

THIR UNITED STATES OF AMIRIOA

Hioneer Hi-Bred International, Inc.

DICCUS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLEMISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE SECULUTE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR ORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'PHADA'

In Testimonn Mercest, I have hereunto set my hand and caused the seal of the Hunt Unriche Frotection Office to be affixed at the City of Washington, D.C. this twenty-third day of November, in the year two thousand and seven.

Attasts

gerzui

Commissioner Plant Variety Protection Offico Agricultural Markoting Service

REPRODUCE LOCALLY, Include form number and	date on all reproc	ductions			Form Approved - OMB No. 0581-0055		
	NT OF AGRICULT		The following statements are made in the Paperwork Reduction Act (PRA)		ce with the Privacy Act of 1974 (5 U.S.C. 552a) and		
SCIENCE AND TECHNOLOGY - F APPLICATION FOR PLANT VA (Instructions and information co	LANT VARIETY PI RIETY PROTECTI	ROTECTION OFFICE ON CERTIFICATE	Application is required in order to dete	rmine if a	plant variety protection certificate is to be issued until certificate is issued (7 U.S.C. 2426).		
NAME OF OWNER			2. TEMPORARY DESIGNATION OR	3. VA	RIETY NAME		
Pioneer Hi-Bred	l Internatio	nal, Inc.	EXPERIMENTAL NAME		PHADA_		
ADDRESS (Street and No., or R.F.D. No., City,	State, and ZIP Co	de, and Country)	5. TELEPHONE (include area code)		FOR OFFICIAL USE ONLY		
7004 1984	62 nd Avenu		515/270-4051	PVPO	NUMBER		
Johnston,			6. FAX (include area code)	1 2	00500223		
comiston,	IA 00 10 1-01	, , , , , , , , , , , , , , , , , , ,	515/253-2125		G DATE		
IF THE OWNER NAMED IS NOT A "PERSON", ORGANIZATION (corporation, partnership, asso		8. IF INCORPORATED, GIVE STATE OF INCORPORATION	9. DATE OF INCORPORATION		April 25,2005		
Corporation		lowa	March 5, 1999	')*		
O. NAME AND ADDRESS OF OWNER REPRES	ENTATIVE(S) TO S	SERVE IN THIS APPLICATION. (First a	erson listed will receive all papers)	F	FILING AND EXAMINATION FEES:		
		·		E S	\$ 3652.00		
		R. Anderson		R E	DATE 4/25/05		
Rese		roduct Development		C E	CERTIFICATION FEE:		
		. Box 85 , IA 50131-0085		V E	\$ 768.00		
				D	DATE 10/15/07		
1. TELEPHONE (Include area code) 515/270-4051	12. FAX (Includ		13. E-MAIL				
4. CROP KIND (Common Name)	16. FAMILY NA	515/253-2125	18. DOES THE VARIETY CONT		Iderson@pioneer.com		
(55,000)		1_233,1031,	☐ YES ☒ NO		(3) 113.11.11		
Corn		Gramineae	ADDDOVED DETITION TO		D USDA-APHIS REFERENCE NUMBER FOR THE LATE THE GENETICALLY MODIFIED PLANT FOR		
5. GENUS AND SPECIES NAME OF CROP	17. IS THE VAR	RIETY A FIRST GENERATION HYBRID	COMMERICALIZATION.	PEREGOI	EATE THE GENETICALET MODIFIED FORTH ON		
Zea Mays							
CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions on reverse)					SEED OF THIS VARIETY BE SOLD AS A CLASS 83(a) of the Plant Variety Protection Act)		
a. ⊠ Exhibit A. Origin and Breeding History of the Variety b. ⊠ Exhibit B. Statement of Distinctness c. ⊠ Exhibit C. Objective Description of Variety d. □ Exhibit D. Additional Description of the Variety (Optional) e. ⊠ Exhibit E. Statement of the Basis of the Owner's Ownership f. ☑ Voucher Sample (2,500 viable untreated seeds or, for tuber propagated varieties,				☐ YES (If "yes", answer items 21 and 22 below) ☑ NO (If "no", go to item 23) 21. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO			
			NUMBER OF CLASSES?				
			☐ YES ☐ NO		• •		
verification that tissue culture will be depo					INDATION ☐ REGISTERED ☐ CERTIFIED SEED OF THIS VARIETY BE LIMITED AS TO		
g. Filing and Examination Fee (\$3,652), made States" (Mail to the Plant Variety Protection		surer of the United	NUMBER OF GENERATION				
			☐ YES ☐ NO				
			IF YES, SPECIFY THE NUM	BER 1,2,3,	, etc. FOR EACH CLASS.		
				EGISTER	ED CERTIFIED please use the space indicated on the reverse.)		
3. HAS THE VARIETY (INCLUDING ANY HARVE			24. IS THE VARIETY OR ANY C	24. IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY INTELLECTUAL PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)?			
FROM THIS VARIETY BEEN SOLD, DISPOSE OTHER COUNTRIES?	D OF, TRANSFER	RED, OR USED IN THE U. S. OR	_				
	TIDOT CALE DICE	OCITION TRANSFER OF LICE	☐ YES ☑ NO				
FOR EACH COUNTRY AND THE CIRCUMSTA	NCES. (Please us	se space indicated on reverse.)	IF YES, PLEASE GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED REFERENCE NUMBER. (Please use space indicated on reverse.)				
The owners declare that a viable sample of bas for a tuber propagated variety a tissue culture v				accordan	ce with such regulations as may be applicable, or		
The undersigned owner(s) is(are) the owner of entitled to protection under the provisions of Se			, and believe(s) that the variety is new,	distinct, un	iform, and stable as required in Section 42, and is		
Owner(s) is (are) informed that false representa	tion herein can jeo	pardize protection and result in penalti	es.				
IGNATURE OF OWNER		SI	GNATURE OF OWNER		0		
			Street.	Ans	leson		
AME (Please print or type)		N	AME (Please print or type)				
			Ste	ven R.	Anderson		
APACITY OR TITLE	DATE	- C/	APACITY OR TITLE Research Scientist	DATE	4-20-2005		
· · · · · · · · · · · · · · · · · · ·							

T 470 (04 03) decianed by the Ellent Mariety Embedden Office union Marie 2003

200500223

3ENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Compléted application form signed by the owner; (2) compléted exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 liling fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. **DO NOT** use masking materials o make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuanc of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvpindex.htm

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that name has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, 10301 Baltimore Avenue, Suite 401 NAL Building, Beltsville, MD 20705. Telephone: (301) 504-5682 http://www.ams.usda.gov/lsg/seed.htm.

TEM

19a.Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively:
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d.Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e.Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

United States, Nov. 1, 2004

24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filing a change of address; owner's representative; ownership or assignment; or any nodification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

coording to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing astructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

he U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, placed by the program of the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, placed by the program of the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, placed by the program of the basis of race, color, national origin, gender, religion, age, disability, sexual orientation in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, placed by the program of the basis of race, color, national origin, gender, religion, age, disability, sexual orientation or family status, placed by the program or family status, p

o file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD) SDA is an equal opportunity provider and employer.

iT-470 (04-03) designed by the Plant Variety Protection Office using Word 2002.

Exhibit A: Developmental history for PHADA

Pedigree: PH38D/PH09B)X12212211X

Pioneer Line PHADA, Zea mays L., a yellow endosperm, dent corn, inbred, was developed by Pioneer Hi-Bred International, Inc. from the single cross hybrid PH38D (PVP certificate Number 9700225) X PH09B (PVP certificate Number 9700218) using the pedigree method of plant breeding. Varieties PH38D and PH09B are proprietary inbred lines of Pioneer Hi-Bred International, Inc. Selfing was practiced from the above hybrid for 10 generations using pedigree selection. During line development, crosses were made to inbred testers for the purpose of estimating the line's combining ability. Yield trials were grown at York, Nebraska as well as other Pioneer research locations. After initial testing, additional hybrid combinations have been evaluated and subsequent generations of the line have been grown and hand-pollinated with observations again made for uniformity.

Variety PHADA has shown uniformity and stability for all traits as described in Exhibit C - "Objective Description of Variety". It has been self-pollinated and ear-rowed 9 generations with careful attention paid to selection criteria and uniformity of plant type to assure genetic homozygousity and phenotypic stability. The line has been increased both by hand and in isolated fields with continued observations for uniformity and stability, and for 7 generations during the final stages of inbred development and seed multiplication. Very high standards for genetic purity have been established morphologically using field observations and using sound lab electrophoresis methodology.

No variant traits have been observed or are expected in PHADA.

The criteria used in the selection of PHADA were yield, both per se and in hybrid combinations. Late season plant health and late season root lodging, grain quality, stalk lodging resistance, and kernel size are especially important in production, and were also important criteria considered during selection. Other selection criteria include: good drought tolerance, brittle snap tolerance, and headsmut tolerance, insect resistance, pollen yield and tassel size.



Exhibit A: Developmental history for PHADA

Pedigree Grown Season/Year	Inbreeding Level of Pedigree Grown
PH38D	F0
РН09В	F0
PH38D/PH09B Winter 1995	F1
PH38D/PH09B)X Summer 1996	F2
PH38D/PH09B)X1 Summer 1997	F3
PH38D/PH09B)X12 Winter 1997	F4
PH38D/PH09B)X122 Summer 1998	F5
PH38D/PH09B)X1221 Summer 1999	F6
PH38D/PH09B)X12212 Winter 1999	F7
PH38D/PH09B)X122122 Summer 2000	F8
PH38D/PH09B)X1221221 Sinter 2000	F9
PH38D/PH09B)X12212211 Summer 2001	F10
PH38D/PH09B)X12212211X	F11 (Seed)

^{*}PH38D was selfed and ear-rowed from F2 through F10 generation. #Uniformity and stability were established from F4 through F10 generation and beyond when seed supplies were increased.

Exhibit B: Novelty Statement

Variety PHADA mostly resembles Pioneer Hi-Bred International, Inc. proprietary inbred line PHP38 (PVP Certificate No. 9000250). Table 1 shows two sample t-tests on data collected primarily in Johnston and Dallas Center, Iowa in 2002 and 2004. The traits collectively show measurable differences between the two varieties.

Exhibit B: Novelty Statement

Variety PHADA has a smaller kernel thickness (4.6 mm vs 6.0 mm) than variety PHP38 (Table 1).

Variety PHADA has a shorter leaf length (74.8 cm vs 85.6 cm) than variety PHP38 (Table 1).

Exhibit B: Novelty Statement Table(s)

ting evidence nments had Table 1: Data from Johnston and Dallas Center, Iowa in 2002 and 2004 presented by trait, across years

for differe different p	ances betwo	een PH Ites and	ADA and were in	for differences between PHADA and PHP38. Each year varieties were grown in 3 locations that had different environmen different planting dates and were in different fields. A two-sample t-test was used to compare differences between means.	Each y	ear vari	eties wample t	ere grow test was	n in 3 loc s used to	uait, ao cations tl compare	hat had differe	differen	oroken (t enviro ween m	out oy nmen teans.	year. D tal cond	for differences between PHADA and PHP38. Each year varieties were grown in 3 locations that had different environmental conditions. Environmental different planting dates and were in different fields. A two-sample t-test was used to compare differences between means.	# #
Kernel thickness (mm)	ess (mm)									· · }							
Level	Station	Year	Year Variety-1	Variety-2	Cnt-1	Cnt-2	Mean-1	Mean-2	Mean-1 Mean-2 Mean_Diff StDev-1 StDev-2 StErr-1 StErr-2 DF	StDev-1	StDev-2	StErr-1	StErr-2	된	t-Value	Prob Pool	
1. Over All			PHADA	PHP38	30	30	4.6	0.9	-14	0.568	0.695	0.104	0.127	58	-8.7	0.00	
2. Year		2002	PHADA	PHP38	15	5	4; 5;	5.9	-1.4	0.516	0.743	0.133	0.192	28	-6.0	0.000	
2. Year		2004	PHADA	PHP38	15	15	4.7	6.1	-1.5	0.617	0.640	0.159	0.165	28	6.4	0.000	

Level Station Year Variety-1 Variety-2 Cnt-1 Cnt-2 Mean-1 Mean-2 Mean-2 Mean-1 StDev-1 StDev-2 StErr-1 StErr-1 DF t-Value Prob_Pool 1. Over All PHADA PHP38 30 74.8 85.6 -10.8 2.833 5.443 0.517 0.994 58 -9.6 0.000 2. Year 2002 PHADA PHP38 15 75.1 85.1 -10.0 3.159 5.139 0.816 1.327 28 -6.4 0.000 2. Year 2. Year 2.004 PHADA PHP38 15 74.5 86.1 -11.6 2.532 5.873 0.654 1.516 28 -7.0 0.000	Lear lengtn (cm	сm)	1							٠						•
All PHADA PHP38 30 74.8 85.6 -10.8 2.833 5.443 0.517 0.994 58 -9.6 2002 PHADA PHP38 15 15 75.1 85.1 -10.0 3.159 5.139 0.816 1.327 28 -6.4 2004 PHADA PHP38 15 15 74.5 86.1 -11.6 2.532 5.873 0.654 1.516 28 -7.0	Level	Station	Year	Variety-1	Variety-2	Cnt-1	Cnt-2	Mean-1			StDev-1	StDev-2	StErr-1	StErr-2	t-Value	Prob Pool
2002 PHADA PHP38 15 15 75.1 85.1 -10.0 3.159 5.139 0.816 1.327 28 -6.4 2004 PHADA PHP38 15 15 74.5 86.1 -11.6 2.532 5.873 0.654 1.516 28 -7.0	1. Over All			PHADA	PHP38	30	30	74.8			2.833	5.443	0.517	0.994	9.6	0.000
2004 PHADA PHP38 15 15 74.5 86.1 -11.6 2.532 5.873 0.654 1.516 28 -7.0	2. Year		2002		PHP38	15	€.	75.1	85.1	-10.0	3.159	5.139	0.816	1.327	-6.4	0.000
	2. Year		2004		PHP38	15	15	74.5	86.1	11.6	2.532	5.873	0.654	1.516	-7.0	0.000

Based on previous discussions with the PVP office the traits longitudinal creases and marginal leaf waves were not collected. These traits have low distinguishing power and are variable due to daily fluctuations in water status of the plants. Therefore, we eliminated them from our process based on previous feedback from the PVP office. For insect or disease traits we included data from disease pressure locations only if they were available and paired with the public check. Most often diseases and insect trials are conducted on hybrids since that is the product ultimately sold. In addition, creating consistent disease pressure and infestation levels is costly and difficult.

In cases where less than 15 observations are presented the trait was collected at the plot level as it always has been done in the past. This means many more plants were visually evaluated according to the procedure outlined below, and then a score of the "population" of the plants was recorded for each location.

The experimental design and methods for 2004 were as follows:

The experiment procedures involved three environments with different planting dates per year, planted in 17.42 ft. rows with 2 rows for each variety. Approximately 24-30 plants emerged in each of 2 rows for a total of around 48 to 60 plants being evaluated at each location and 144 to 180 plants across locations. For plant level traits, we sampled 5 representative plants from the 2 rows of the 2 row plot (group) of plants at each location. For plot level traits we evaluated the 2 row plot (group) and gave a representative score or average on the 48-60 plants in the group within an experiment.

Some traits can be especially variable under different environmental factors influenced by weather, soil type, or planting dates. Varying temperatures or day length could impact the meristem growth during various tissue differentiation stages. The meristem differentiation of the ear and other tissues could be impacted as well as the success of pollination during flowering and frequency of kernel abortion during grain fill.

We have included weather data in the table that follows.

Month	GROW	GROWING DEGREE UNITS (GDU'S)	EE UNITS (GDU's)	ā	PRECIPITATION	ION (inches)	19
	20	2003	20	2004	20	2003		2004
	D. Center	Johnston	D. Center	Johnston	D. Center	Johnston	D. Center	Johnston
May	375	380	548	527	5.7	5.43	7 19	7.75
June	909	604	609	610	1.92	4.23	1 97	3 30
July	628	782	723	736	0.18	3.4	2 29	4 54
August	795	786	612	615	0.44	0.51	1.95	4 95
September	456	468	598	560	2.19	2.52	138	1 24
TOTAL	2860	3020	3090	3048	10.43	16.09	14.78	21.87

Calculate GDU's

Growing Degree Units use following formula: GDU = ((T1+T2)/2)-50

Where T2 = maximum temperature for a given day with 86 degrees Fahrenheit as the maximum temperature used and 50 degrees Fahrenheit is the minimum temperature used. Where T1 = minimum temperature for a given day with 50 degrees Fahrenheit as the minimum temperature used and 86 degrees Fahrenheit is the maximum temperature used. GDU's are calculated each day and accumulated (summed) over certain number of days.

United States Department of Agriculture, Agricultural Marketing Service Science and Technology, Plant Variety Protection Office National Agricultural Library Building, Room 400 Beltsville, MD 20705-2351

OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

Name of Applicant(s		l Variety Seed	Source		Temporary Designation
Pioneer Hi-Bred Int	ernational, inc	1		1 PHADA	
Address (Street & No 7301 NW 62nd Aver	o., or R.F.D. No., City, State, nue, P.O. Box 85, Johnston	Zip Code and Country , lowa 50131-0085	I FOR OFFICIA	1 use 2 0	PVPO Number 0 5 0 0 2 2 3
adding leading zeroe	e number that describes the es if necessary. Completenes ry for an adequate variety des	s should be striven for to	establish an adequate va	ne spaces below. Right riety description. Traits	justify whole numbers by designated by a "*" are
COLOR CHOICES (I 01. Light Green 02. Medium Green 03. Dark Green 04. Very Dark Gree 05. Green-Yellow	Use in conjunction with Muns 06. Pale Yellow 07. Yellow 08. Yellow-Orange n 09. Salmon 10. Pink-Orange	ell color code to describe 11. Pink 12. Light Red 13. Cherry Red 14. Red 15. Red & White	all color choices; descrit 16. Pale Purple 17. Purple 18. Colorless 19. White 20. White Capped	pe #25 and #26 in Comr 21. Buff 22. Tan 23. Brown 24. Bronze 25. Variegated (Des	26. Other (Describe)
Yellow Dent Familie Family B14 B37 B73 C103 Oh43	D CHOICES [Use the most si es: Members CM105, A632, B64, B68 B37, B76, H84 N192, A679, B73, Nc268 Mo17, Va102, Va35, A682 A619, MS71, H99, Va26 W64A, A554, A654, Pa91	milar (in background and Yellow Dent (Unrelated Co109, ND246 Oh7, T232 W117, W153R W182BN White Dent: Cl66, H105, Ky2):	Sweet Corn: C13, Iowa512 Popcorn:	25, P39, 2132 722, HP301, HP7211
	intermediate types in "Comm tt, 2=Dent, 3=Flint, 4=Flour, 5 t		Pipecorn)	I Standard Inbred I 2 Type	
	E DEVELOPED IN THE U.S./ est, 2=N.Central, 3=N.East, 4	and the second s	S.West, 7=Other	I Standard Seed S	ource AMES 19306
	egion Best Adaptability, show HEAT UNITS 1,486.4 From emergence 1,468.7 From 10% to 90% From 50% silk to	to 50% of plants in silk to 50% of plants in polle	n	DAYS 57 56 2	HEAT UNITS 1,289.7 1,258.7 52
86.7 cm Ear I 15.8 cm Leng 0.0 Average 1.2 Average	Height (to tassel tip) Height (to base of top ear noo th of Top Ear Internode Number of Tillers Number of Ears per Stalk anin of Brace Roots: 1=Abse	te)	15.45 2.37 0.01 0.27	193.5 1 51.6 1 15.2 1 0.0	St.Dev. Sample Size 15.23 45 8.24 45 1.73 45 0.02 9 0.07 9
Application Variety D	Data		Page 1	I Standard Inbred	Data

pplication Variety Data	Page 2		. I Sta			
LEAF		0	an action of a 162 or decode will refer	5.7	04.5	
10.1 cm Width of Ear Node Leaf	St.Dev.	Sample Siz		Mean	St.Dev.	Sample S
	<u>0.89</u>		<u>5</u>	<u>9.5</u>	0.97	
75.8 cm Length of Ear Node Leaf	4.22	4	<u>5</u> 1	<u>68.4</u>	<u>5.21</u>	
6.8 Number of leaves above top ear	<u>0.71</u>		<u>5</u> I	<u>5.4</u>	<u>0.65</u>	
21.7 Degrees Leaf Angle	<u>3.84</u>	4	<u>.5</u> l	<u>25.4</u>	7.42	
(Measure from 2nd leaf above ear at anthesis to stalk about	ove leaf)	•	1			
4 Leaf Color (Munsell Code) 5GY34			I .	4 (Munsell	Code) 5G	Y34
5 Leaf Sheath Pubescence (Rate on scale from 1=none to	9=like peach	fuzz)	· 1	5	, <u></u>	
Marginal Waves (Rate on scale from 1=none to 9=many	y (- + + + + + + + + + + + + + + + + + +	,,	i	_		
Longitudinal Creases (Rate on scale from 1=none to 9=	many)		i :		*	
						· · · · · · · · · · · · · · · · · · ·
TASSEL:	St.Dev.	Sample Siz		Mean	St.Dev.	Sample Si
4.6 Number of Primary Lateral Branches	<u>1.66</u>	. <u>4</u>	<u>5</u> l	<u>10.5</u>	<u>2.36</u>	
20.6 Degrees Branch Angle from Central Spike	<u>7.91</u>	4	<u>5</u>	<u>29.7</u>	<u>7.77</u>	
56.8 cm tassel Length	<u>3.18</u>	. 4	<u>5</u> 1	<u>58.2</u>	4.76	
(from top leaf collar to tassel tip)			· [
5 Pollen Shed (Rate on scale from 0=male sterile to 9=he	avy shed)		I	7		
12 Anther Color (Munsell Code) 7.5R58				8 (Munsell	Code) <u>10Y</u>	8/10
14 Glume Color (Munsell Code) 2.5R46			Ī		Code) <u>5G</u>	
1 Bar Glumes (Glume Bands): 1=Absent, 2=Present			i ·	1		72
EAR (Unhusked Data):				_		
<u>12</u> Silk Color (3 days after emergence) (Munsell Code)	5R:	sk	· I	1 Munsell (Code 250	GY9 4
2 Fresh Husk Color (25 days after 50% silking) (Munsell (8	1	1 Munsell (
2 Treen trues Color (25 days after 50% Sliking) (Munsell C	do) 100		1			
19 Dry Husk Color (65 days after 50% silking) (Munsell Co		<u>(R92</u>	i	21 Munsell	Code <u>2.5</u>	<u>18.5</u> 4
1 Position of Ear at Dry Husk Stage: 1=Upright, 2=Horizo	ntal 3≍Penden	4 (-		5
		t ·	!	2		. 1
6 Husk Tightness (Rate on scale from 1=very loose to 9=	very tight			· <u>∠</u> <u>6</u>		. 1
2 Husk Extension (at harvest): 1=Short(ears exposed), 2=	very tight		 	<u>∠</u> <u>6</u> <u>3</u>		
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm)	very tight ∍Medium (<8cm	n), 3=Long		<u>6</u> <u>3</u>		. :
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data)	very tight •Medium (<8cm St. Dev.	n), 3=Long Sample Siz		6 3 Mean	St.Dev.	Sample Si
 2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 	very tight =Medium (<8cm St. Dev. 1.37	n), 3=Long Sample Siz 4	5 1	6 3 Mean 14.0	<u>2.04</u>	
 2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 	very tight -Medium (<8cm St. Dev. 1.37 3.45	Sample Siz	<u>5</u> l <u>5</u> l	6 3 Mean 14.0 45.4	2.04 2.48	
 2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 	very tight =Medium (<8cm St. Dev. 1.37	Sample Siz	<u>5</u> <u>5</u> 5	6 3 Mean 14.0	<u>2.04</u>	
 2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 	very tight -Medium (<8cm St. Dev. 1.37 3.45	Sample Siz	<u>5</u> <u>5</u> 5	6 3 Mean 14.0 45.4 95.0	2.04 2.48	
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30	Sample Siz	<u>5</u> <u>5</u> 5	Mean 14.0 45.4 95.0 14.8	2.04 2.48 21.34	Sample Si
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct	very tight •Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30	Sample Siz	<u>5</u> <u>5</u> 5	Mean 14.0 45.4 95.0 14.8 2	2.04 2.48 21.34	
 2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 	very tight •Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30	Sample Siz	5 5 5 5 1	Mean 14.0 45.4 95.0 14.8 2	2.04 2.48 21.34 1.57	
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length	very tight •Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73	Sample Siz	5 5 5 5 1	Mean 14.0 45.4 95.0 14.8 2	2.04 2.48 21.34	
 2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 	very tight •Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73	Sample Siz	5 5 5 5 1	Mean 14.0 45.4 95.0 14.8 2	2.04 2.48 21.34 1.57	
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried):	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St.Dev.	Sample Siz 4 4 4 Sample Siz	5 5 5 5 5 5 5 5 5 5	Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean	2.04 2.48 21.34 1.57 3.10 St.Dev.	
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St. Dev. 1.10	Sample Siz 4 4 4 Sample Siz	5 5 5 5 5 5 5 5 5 5	Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92	
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St.Dev. 1.10 0.69	Sample Siz Sample Siz 4 4 4 Sample Siz	55 1 1 1 1 1 1 1 1 1	Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77	Sample S
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Thickness	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St. Dev. 1.10 0.69 0.65	Sample Siz 4 4 Sample Siz 4 4 Sample Siz		Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86	
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade)	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St.Dev. 1.10 0.69 0.65 15.09	Sample Siz 4 4 Sample Siz 4 4 Sample Siz	55 1 1 1 1 1 1 1 1 1	Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77	
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St.Dev. 1.10 0.69 0.65 15.09	Sample Siz 4 4 Sample Siz 4 4 Sample Siz		Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13	
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade)	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St.Dev. 1.10 0.69 0.65 15.09 (describe)	Sample Siz 4 4 Sample Siz 4 4 Sample Siz		Mean 14.0 45.4 95.0 14.8 2 12.4 2 10.9 8.6 5.1 47.4	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13	
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Length 4.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St.Dev. 1.10 0.69 0.65 15.09 (describe)	Sample Siz 4 4 Sample Siz 4 4 Sample Siz		Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1 47.4 1 (describe	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13 e)	Sample S
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating 7 Aleurone Color (Munsell Code) 1 Hard Endosperm Color (Munsell Code)	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St. Dev. 1.10 0.69 0.65 15.09 (describe) 14 12	Sample Siz Sample Siz 4 4 Sample Siz 4 Sample Siz		Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1 47.4 1 (describe 7 Munsell (7 Mun	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13 St.Dev. 15.13	Sample S
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating 7 Aleurone Color (Munsell Code) 1 Hard Endosperm Color (Munsell Code) 3 Endosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St. Dev. 1.10 0.69 0.65 15.09 (describe) 14 12 3=Normal Starc	Sample Siz Sample Siz 4 4 Sample Siz 4 Sample Siz 4 4 h, 4=High		Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1 47.4 (describe 7 Munsell (2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13 St.Dev. 15.13	Sample S
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Hornozygous, 2=Segregating 7 Aleurone Color (Munsell Code) 1 10YR7 7 Hard Endosperm Color (Munsell Code)	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St. Dev. 1.10 0.69 0.65 15.09 (describe) 14 12 3=Normal Starc	Sample Siz Sample Siz 4 4 Sample Siz 4 Sample Siz 4 4 h, 4=High		Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1 47.4 1 (describe 7 Munsell (7 Mun	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13 St.Dev. 15.13	Sample S
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating 7 Aleurone Color (Munsell Code) 1 Aleurone Color (Munsell Code) 2 Endosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3 Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High (se), 9=High Oil, 10=Other	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St. Dev. 1.10 0.69 0.65 15.09 (describe) 14 12 =Normal Starce Lysine, 8=Sup	Sample Siz 4 4 Sample Siz 4 4 Sample Siz 4 4 And	55 1 1 1 1 1 1 1 1 1	Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1 47.4 1 (describe 7 Munsell (7 Munsell (3 (describe	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13 Code 1	Sample S
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating 7 Aleurone Color (Munsell Code) 1 Aleurone Color (Munsell Code) 3 Endosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3 Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High (se), 9=High Oil, 10=Other 28.1 gm Weight per 100 kernels (unsized sample)	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St. Dev. 1.10 0.69 0.65 15.09 (describe) 14 12 3=Normal Starc	Sample Siz 4 4 Sample Siz 4 4 Sample Siz 4 4 And		Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1 47.4 1 (describe 7 Munsell (7 Mun	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13 St.Dev. 15.13	Sample S
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating 7 Aleurone Color (Munsell Code) 1 Aleurone Color (Munsell Code) 3 Endosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3 Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High (se), 9=High Oil, 10=Other 28.1 gm Weight per 100 kernels (unsized sample)	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St. Dev. 1.10 0.69 0.65 15.09 (describe) 14 12 =Normal Starce Lysine, 8=Sup	Sample Siz 4 4 Sample Siz 4 4 Sample Siz 4 4 And		Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1 47.4 1 (describe 7 Munsell (7 Munsell (3 (describe	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13 Code 1 2.00de 1 3.33	Sample S 0YR8[14 0YR7[12
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating 7 Aleurone Color (Munsell Code) 1 Aleurone Color (Munsell Code) 2 Endosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3 Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High (se), 9=High Oil, 10=Other 28.1 gm Weight per 100 kernels (unsized sample) OB: 24.8 mm Cob Diameter at mid-point	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St.Dev. 1.10 0.69 0.65 15.09 (describe) 14 12 =Normal Starc Lysine, 8=Sup	Sample Siz 4 4 4 Sample Siz 4 4 Sample Siz 4 4 4 Sample Siz 4 4 4 Sample Siz 4 4 4 4 4 Sample Siz 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	55 55	Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1 47.4 1 (describe 7 Munsell (describe) 2 (describe)	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13 Elode 1 1	Sample S
2 Husk Extension (at harvest): 1=Short(ears exposed), 2= (8-10cm beyond ear tip), 4=Very Long (>10cm) EAR (Husked Ear Data) 15.3 cm Ear Length 43.8 mm Ear Diameter at mid-point 133.0 gm Ear Weight 17.2 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 9.0 cm Shank Length 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Ext [ERNEL (Dried): 11.7 mm Kernel Length 7.6 mm Kernel Width 4.6 mm Kernel Thickness 41.8 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating 7 Aleurone Color (Munsell Code) 1 Aleurone Color (Munsell Code) 3 Endosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3 Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High (se), 9=High Oil, 10=Other 28.1 gm Weight per 100 kernels (unsized sample) OB:	very tight -Medium (<8cm St. Dev. 1.37 3.45 35.30 1.30 1.73 reme conical St.Dev. 1.10 0.69 0.65 15.09 (describe) 14 12	Sample Siz 4 4 4 Sample Siz 4 4 Sample Siz 4 4 5 h, 4=High her Sweet	55 55	Mean 14.0 45.4 95.0 14.8 2 12.4 2 Mean 10.9 8.6 5.1 47.4 1 (describe 7 Munsell (7 Munsell (3 (describe	2.04 2.48 21.34 1.57 3.10 St.Dev. 0.92 0.77 0.86 15.13 St.Dev 2.50de 1 1 St.Dev 2.51	Sample S 0YR8[14 0YR7[12

2 00 5 0022 3 Standard Inbred Data

10. DISEASE RESISTANCE (Rate from 1(most susceptible) to 9 (i	most resistant): lea	ave blank		
if not tested; leave Race or Strain Options blank if polygenic):				e e e
A. Leaf Blights, Wilts, and Local Infection Diseases				·
_ Anthracnose Leaf Blight (Colletotrichum graminicola)			Anthracnose Leaf Blight	
Common Rust (Puccinia sorghi)			I Common Rust	
_ Common Smut (Ustilago maydis)		•	I Common Smut	
Eyespot (Kabatiella zeae)			I Eyespot	
Goss's Wilt (Clavibacter michiganense spp. nebraskensis)			Goss's Wilt	
5 Gray Leaf Spot (Cercospora zeae-maydis)			3 Gray Leaf Spot	
Helminthosporium Leaf Spot (Bipolaris zeicola)	Race		Helminthosporium Leaf Spot	Race
6 Northern Leaf Blight (Exserohilum turcicum)	Race	·	I 3 Northern Leaf Blight	Race
Southern Leaf Blight (Bipolaris maydis)	Race		I Southern Leaf Blight	Race_
Southern Rust (Puccinia Polysora)			I Southern Rust	11000
Stewart's Wilt (Erwinia stewartii)			I Stewart's Wilt	
Other (Specify)			Other (Specify)	
B. Systemic Diseases				_
Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MDMV) Strain Sorghum Downy Mildew of Corn (Peronosclerospora sorg Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme)	hi)	<u>-</u>	Corn Lethal Necrosis Head Smut Maize Chlorotic Dwarf Virus Maize Chlorotic Mottle Virus Maize Dwarf Mosaic Virus Sorghum Downy Mildew of C Other (Specify) Anthracnose Stalk Rot Diplodia Stalk Rot Fusarium Stalk Rot	Strain corn
Gibberella Stalk Rot (Gibberella zeae) Other (Specify)			Gibberella Stalk Rot	
D. Ear and Kernel Rots			_ Other (Specify)	 -
— · · · · · · · · · · · · · · · · · · ·	50.0	•		
 Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarim Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) 			I Aspergillus Ear & Kernel Rot I Diplodia Ear Rot I Fusarium Ear & Kernel Rot I Gibberella Ear Rot	
_ Other (Specify)			I Other (Specify)	
Application Variety Data	Page 3		I Standard Inbred Data	<u></u>

Note: Use chart on first page to choose color codes for color traits.

	an yanning mana arawa animana a minajanina ming agawa najaaj njigganga aga	· 		1Standard Inbred Data	enderstanding and control of a more
11. INSECT RESISTAN	CE (Rate from 1(most susceptib			1	
if not tested	er en	St. Dev.	Sample Size	I St. Dev.	Sample Si
	lite (Oligonychus pratensis)			Banks Grass Mite	
Corn Earworm (Helio	coverpa zea)		4.	Corn Earworm	
_ Leaf Feeding				Leaf Feeding	•
	mg larval wt.				
_ Ear Damage	d (Dhanalasinis (m. m. sidis)			_ Ear Damage	
	d (Rhopalosiphum maidis)		·	Corn Leaf Aphid	
	le (Carpophilus dimidiatus)			Corn Sap Beetle	
European Corn Bore				European Corn Borer	
	(Typically Whorl Leaf Feeding)			1 st Generation	
_ 2 nd Generano	n (Typically Leaf Sheath-Collar F	-eeaing)		_ 2 nd Generation	
	cm tunneled/plant	·			
Fall Armyworm (Spor	doptera frugiperda)			Fall Armyworm	
_ Leaf-Feeding				_ Leaf-Feeding	-
	mg larval wt.				
_ iviaize vveevii (3	Sitophilus zeamais)			Maize Weevil	
	rorm (Diabrotica barberi)			_ Northern Rootworm	
	vorm (Diabrotica undecimpuncta	ta)	1	Southern Rootworm	
Southwestern Corn B	Borer (Diatraea grandiosella)		· I	Southwestern Corn Borer	
_ Leaf Feeding			I	_ Leaf Feeding	
Stalk Tunneling:	cm tunneled/plant				
Two-spotted Sp	ider Mite (Tetranychus urticae)	· · · · · · · · · · · · · · · · · · ·	· · · · · · · · · · · · · · · · · ·	Two-spotted Spider Mite	
_ Western Rootwe	orm (Diabrotica virgifera virgifera	a) .		Western Rootworm	
<pre>_ Other (Specify)</pre>			I	_ Other (Specify)	
				<u> </u>	_
% Dropped Ears _ % Pre-anthesis 0 % Pre-anthesis 0 % Post-anthesis	65 days after anthesis) (Rate on (at 65 days after anthesis) Brittle Snapping	anthesis)	9=excellent) [Stay Green Dropped ears Pre-anthesis Brittle Sna Pre-anthesis Root Lodg Post-anthesis Root Lodgir 3.835.0 Yield 	ing
13. MOLECULAR MARKI	ERS: (0=data unavailable; 1=da	ta available but not su	pplied; 2=data supplied	<u>. </u>	
4					
1 Isozymes	_ RFLP's	_ RAPD's		Other (Specify)	-
. '					
REFERENCES:			•		
Emerson, R.A., G.W. Bea	em for the Classification of Corn adle, and A.C. Fraser, 1935. A s . Chamuris, A.Y. Rossman. 198	ummary of Linkage S	udies in Maize. Cornell	I A.E.S., Mem. 180. tes. The American Phytopathological	
Inglett, G.E. (Ed) 1970. C Jugenheimer, R.W. 1976. McGee, D.C. 1988. Maize Munsell Color Chart for P The Mutants of Maize. 19 Shurtleff, M.C. 1980. Com	orn: Culture, Processing, Product. Corn: Improvement, Seed Product Diseases. APS Press, St. Paul lant Tissues. Macbeth. P.O. Box 68. Crop Science Society of Ampendium of Corn Diseases. APS Dudley (Editors). 1988. Corn and	luction, and Uses. Jol , MN. 150 pp. (230. Newburgh, N.Y erica. Madison, WI. S Press, St. Paul, MN	n Wiley & Sons, New 12551-0230 . 105 pp.	York. ny Monograph 18. ASA, CSSA, SSSA,	
	ored Lines of Ohio A.E.S., Bul. 8 ulture 1936, 1937. Yearbook.	31. 1959.			

COMMENTS (e.g. state how heat units were calculated, standard inbred seed source, and/or where data was collected. Continue in Exhibit D)
Insect, disease, brittle snapping and root lodging data are collected mainly from environment where variability for the trait can be obtained within the experiment.

CLARIFICATION OF DATA IN EXHIBITS B AND C

Please note the data presented in Exhibit B and C, "Objective Description of Variety," are collected primarily at Johnston and Dallas Center, Iowa. The data in Table 1 are from two sample t-tests using data collected in Johnston and Dallas Center, IA. These traits in Exhibit B collectively show distinct differences between the two varieties.

REPRODUCE LOCALLY. Include form number and edition date on all reproductions.	FORM APPROVED - OMB NO	. 0581-0055
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to detect certificate is to be issued (7 U.S.C. 24 confidential until the certificate is issued	21). The information is held
1.NAME OF APPLICANT(S)	2.TEMPORARY DESIGNATION	3. VARIETY NAME
PIONEER HI-BRED INTERNATIONAL, INC.	OR EXPERIMENTAL NUMBER	PHADA
4 .ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5.TELEPHONE (include area code)	6. FAX (include area code)
7301 NW 62 nd AVENUE	515-270-4051	515-253-2125
P.O.BOX 85	7.PVPO NUMBER	
JOHNSTON, IA 50131-0085	2	00500223
8. Does the applicant own all rights to the variety? Mark an "X" in the appropriate bit	ock. If no, please explain: 🛛 YE	S NO
		• .
9. Is the applicant (individual or company) a U.S. national or a U.S. based company	? If no, give name of country.	YES NO
10. Is the applicant the original owner?	swer <u>one</u> of the following:	
a. If the original rights to variety were owned by individual(s), is (are) the orig	inal owner(s) a U.S. National(s)?	
	,	
☐ YES ☐ NO If no, give name of country		
b. If the original rights to variety were owned by a company(ies), is (are) the	original owner(s) a U.S. based compa	any?
☑ YES ☐ NO If no, give name of country		
11. Additional explanation on ownership (Trace ownership from original breeder to o	current owner. Use the reverse for ex	tra space if needed):
Pioneer Hi-Bred International, Inc. (PHI), Des Moines, Iowa, and/or its wholly is the employer of the plant breeders involved in the selection and developme Corporation has the sole rights and ownership of PHADA pursuant to written out such variety was created. No rights to this variety are retained by any individual.	nt of PHADA. Pioneer Hi-Bred Intern contracts that assign all rights in the v	ational and/or Pioneer Overseas
PLEASE NOTE:		
Plant variety protection can only be afforded to the owners (not licensees) who meet the fo	llowing criteria:	
 If the rights to the variety are owned by the original breeder, that person must be a U which affords similar protection to nationals of the U.S. for the same genus and speci 		er country, or national of a country
If the rights to the variety are owned by the company which employed the original br member country, or owned by nationals of a country which affords similar protection		
3. If the applicant is an owner who is not the original owner, both the original owner an	d the applicant must meet one of the abo	ve criteria.
The original breeder/owner may be the individual or company who directed the final breed	ing. See Section 41(a)(2) of the Plant V	ariety Protection Act for definitions.
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is	not required to respond to a collection of informat	ion unless it displays a valid OMB control number.

reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal employment opportunity provide and employer.